

King's Research Portal

DOI:

[10.1038/s41588-018-0059-2](https://doi.org/10.1038/s41588-018-0059-2)

Document Version

Peer reviewed version

[Link to publication record in King's Research Portal](#)

Citation for published version (APA):

Pardiñas, A. F., Holmans, P., Pocklington, A. J., Escott-Price, V., Ripke, S., Carrera, N., Legge, S. E., Bishop, S., Cameron, D., Hamshere, M. L., Han, J., Hubbard, L., Lynham, A., Mantripragada, K., Rees, E., MacCabe, J. H., McCarroll, S. A., Baune, B. T., Breen, G., ... GERAD1 Consortium: (2018). Common schizophrenia alleles are enriched in mutation-intolerant genes and in regions under strong background selection. *Nature Genetics*, 50, 381–389. <https://doi.org/10.1038/s41588-018-0059-2>

Citing this paper

Please note that where the full-text provided on King's Research Portal is the Author Accepted Manuscript or Post-Print version this may differ from the final Published version. If citing, it is advised that you check and use the publisher's definitive version for pagination, volume/issue, and date of publication details. And where the final published version is provided on the Research Portal, if citing you are again advised to check the publisher's website for any subsequent corrections.

General rights

Copyright and moral rights for the publications made accessible in the Research Portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognize and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the Research Portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the Research Portal

Take down policy

If you believe that this document breaches copyright please contact librarypure@kcl.ac.uk providing details, and we will remove access to the work immediately and investigate your claim.

Common schizophrenia alleles are enriched in mutation-intolerant genes and in regions under strong background selection

Antonio F. Pardiñas^a, Peter Holmans^a, Andrew J. Pocklington^a, Valentina Escott-Price^a, Stephan Ripke^{b,c}, Noa Carrera^a, Sophie E. Legge^a, Sophie Bishop^a, Darren Cameron^a, Marian L. Hamshire^a, Jun Han^a, Leon Hubbard^a, Amy Lynham^a, Kiran Mantripragada^a, Elliott Rees^a, James H. MacCabe^d, Steven A. McCarroll^e, Bernhard T. Baune^f, Gerome Breen^{g,h}, Enda M. Byrneⁱ, Udo Dannlowski^j, Thalia C. Eley^g, Caroline Hayward^l, Nicholas G. Martin^{m,n}, Andrew M. McIntosh^{o,p}, Robert Plomin^g, David J. Porteous^l, Naomi R. Wrayⁱ, Armando Caballero^q, Daniel H. Geschwind^r, Laura M. Huckins^s, Douglas M. Ruderfer^s, Enrique Santiago^t, Pamela Sklar^s, Eli A. Stahl^s, Hyejung Won^r, Esben Agerbo^{u,v,w}, Ole A. Andreassen^x, Marie Bækvad-Hansen^{u,y}, Preben Bo Mortensen^{u,v,w,z}, Carsten Bøcker Pedersen^{u,v,w}, Anders D. Børglum^{u,z,aa}, Jonas Bybjerg-Grauholm^{u,y}, Thomas D. Als^{u,z,aa}, Srdjan Djurovic^{ab}, Naser Durmishi^{ac}, Marianne Giørtz Pedersen^{u,v,w}, Vera Golimbet^{ad}, Jakob Grove^{u,z,aa,ad}, David M. Hougaard^{u,y}, Manuel Mattheisen^{u,z,aa}, Espen Molden^{af}, Ole Mors^{u,ag}, Merete Nordentoft^{u,ah}, Milica Pejovic-Milovancevic^{ai}, Engilbert Sigurdsson^{aj}, Teimuraz Silagadze^{ak}, Christine Søholm Hansen^{u,y}, Kari Stefansson^{al}, Hreinn Stefansson^{al}, Stacy Steinberg^{al}, Sarah Tosato^{am}, Thomas Werge^{u,an,ao}, GERAD1 Consortium^{ap}, David A. Collier^{g,aq}, Dan Rujescu^{ar,as}, George Kirov^a, Michael J. Owen^{a*}, Michael C. O'Donovan^{a*}, James T. R. Walters^{a*}

^a MRC Centre for Neuropsychiatric Genetics and Genomics, Institute of Psychological Medicine and Clinical Neurosciences, School of Medicine, Cardiff University, Cardiff, UK

^b Analytic and Translational Genetics Unit, Massachusetts General Hospital, Boston, Massachusetts, USA

^c Department of Psychiatry and Psychotherapy, Charité, Campus Mitte, 10117 Berlin, Germany

^d Department of Psychosis Studies, Institute of Psychiatry Psychology and Neuroscience, King's College London

^e Stanley Center for Psychiatric Research, Broad Institute of MIT and Harvard, Cambridge, Massachusetts, USA

^f Discipline of Psychiatry, University of Adelaide, Australia

^g Medical Research Council, Social, Genetic and Developmental Psychiatry Centre, Institute of Psychiatry, Psychology and Neuroscience, King's College London, London, UK

^h NIHR Biomedical Research Centre for Mental Health, Maudsley Hospital and Institute of Psychiatry, Psychology and Neuroscience, King's College London, London, UK

ⁱ Queensland Brain Institute, University of Queensland, Brisbane, Queensland, Australia

^j Department of Psychiatry and Psychotherapy University of Muenster, Muenster, Germany

^l Medical Genetics Section, Centre for Genomic and Experimental Medicine, Institute of Genetics and Molecular Medicine, University of Edinburgh, Edinburgh, UK

^m School of Psychology, University of Queensland

ⁿ QIMR Berghofer Medical Research Institute, Brisbane, Queensland, Australia

^o Division of Psychiatry, University of Edinburgh, Edinburgh, UK

^p Centre for Cognitive Ageing and Cognitive Epidemiology, University of Edinburgh, Edinburgh, UK

^q Departamento de Bioquímica, Genética e Inmunología. Facultad de Biología. Universidad de Vigo. Vigo, Spain

^r Department of Neurology, Center for Autism Research and Treatment, Semel Institute, David Geffen School of Medicine, University of California Los Angeles, California, USA.

^s Division of Psychiatric Genomics, Department of Psychiatry, Icahn School of Medicine at Mount Sinai, New York, New York, USA.

^t Departamento de Biología Funcional. Facultad de Biología. Universidad de Oviedo. Oviedo, Spain

^u iPSYCH, The Lundbeck Foundation Initiative for Integrative Psychiatric Research, Denmark

^v National Centre for Register-Based Research, Aarhus University, Aarhus, Denmark

^w Centre for Integrated Register-based Research, Aarhus University, Aarhus, Denmark

^x Institute of Clinical Medicine, University of Oslo, Oslo, Norway

^y Center for Neonatal Screening, Department for Congenital Disorders, Statens Serum Institut, Copenhagen, Denmark

^z iSEQ, Center for Integrative Sequencing, Aarhus University, Aarhus, Denmark

^{aa} Department of Biomedicine - Human Genetics, Aarhus University, Aarhus, Denmark

^{ab} NORMENT, KG Jebsen Centre for Psychosis Research, Department of Clinical Science, University of Bergen, Bergen, Norway

^{ac} Department of Child and Adolescent Psychiatry, University Clinic of Psychiatry, Skopje, Republic of Macedonia

^{ad} Department of Clinical Genetics. Mental Health Research Center, Russian Academy of Medical Sciences. Kashirskoe shosse 34. 115522. Moscow, Russia

^{ae} Bioinformatics Research Centre, Aarhus University, Aarhus, Denmark

^{af} Center for Psychopharmacology, Diakonhjemmet Hospital, Oslo, Norway

^{ag} Psychosis Research Unit, Aarhus University Hospital, Risskov, Denmark

^{ah} Mental Health Services in the Capital Region of Denmark, Mental Health Center Copenhagen, University of Copenhagen, Copenhagen, Denmark

^{ai} Department of Psychiatry, School of Medicine, University of Belgrade, Belgrade, Serbia

^{aj} Department of Psychiatry, National University Hospital, Reykjavik, Iceland

^{ak} Department of Psychiatry and Drug Addiction, Tbilisi State Medical University (TSMU), Tbilisi, Georgia

^{al} deCODE genetics, Reykjavik, Iceland

^{am} Section of Psychiatry, Department of Public Health and Community Medicine, University of Verona, Verona, Italy

^{an} Institute of Biological Psychiatry, MHC Sct. Hans, Mental Health Services Copenhagen, Roskilde, Denmark

^{ao} Department of Clinical Medicine, University of Copenhagen, Copenhagen, Denmark

^{ap} Control data used in the preparation of this article were obtained from the Genetic and Environmental Risk for Alzheimer's disease (GERAD1) Consortium. As such, the investigators within the GERAD1 consortia contributed to the design and implementation of GERAD1 and/or provided control data but did not participate in analysis or writing of this report.

^{aq} Discovery Neuroscience Research, Eli Lilly and Company Ltd, Lilly Research Laboratories, Erl Wood Manor, Surrey, UK

^{ar} Department of Psychiatry, University of Halle, Halle, Germany

^{as} Department of Psychiatry, University of Munich, Munich, Germany

* Correspondence and material requests should be addressed to M.J.O. (owenmj@cardiff.ac.uk), M.C.O'D. (odonovanmc@cardiff.ac.uk) or J.T.R.W. (waltersjt@cardiff.ac.uk)

ABSTRACT

Schizophrenia is a debilitating psychiatric condition often associated with poor quality of life and decreased life expectancy. Lack of progress in improving treatment outcomes has been attributed to limited knowledge of the underlying biology, although large-scale genomic studies have begun to provide such insight. We report a new genome-wide association study of schizophrenia (11,260 cases and 24,542 controls) and through meta-analysis with existing data we identify 50 novel associated loci and 145 loci in total. Through integrating genomic fine-mapping with brain expression and chromosome conformation data, we identify candidate causal genes within 33 loci. We show for the first time that the common variant association signal is highly enriched among genes that are under strong selective pressures. These findings provide novel insights into the biology and genetic architecture of schizophrenia, highlight the importance of mutation intolerant genes and suggest a mechanism by which common risk variants persist in the population.

Schizophrenia is characterised by psychosis and negative symptoms such as social and emotional withdrawal. While onset of psychosis typically does not occur until late adolescence or early adult life, there is strong evidence from clinical and epidemiological studies that schizophrenia reflects a disturbance of neurodevelopment¹. It confers substantial mortality and morbidity, with a mean reduction in life expectancy of 15-30 years^{2,3}. Although recovery is possible, most patients have poor social and functional outcomes⁴. No substantial improvements in outcomes have emerged since the advent of antipsychotic medication in the mid-20th century, a fact that has been attributed to a lack of knowledge of pathophysiology¹.

Schizophrenia is both highly heritable and polygenic, with risk ascribed to variants spanning the full spectrum of population frequencies⁵⁻⁷. The relative contributions of alleles of various frequencies is not fully resolved, but recent studies estimate that common alleles, captured by genome-wide association study (GWAS) arrays, capture between a third and a half of the genetic variance in liability⁸. There has been a long-standing debate, from an evolutionary standpoint, as to how common risk alleles persist in the population, particularly given the early mortality and decreased fecundity associated with schizophrenia⁹. Various hypotheses have been proposed including compensatory advantage (balancing selection), whereby schizophrenia alleles confer reproductive advantages in particular contexts^{10,11}; hitchhiking, whereby risk alleles are maintained by their linkage to positively selected alleles¹²; or contrasting theories that attribute these effects to rare variants and gene-environment interaction¹³. Addressing these competing hypotheses is now tractable given advances from recent studies of common genetic variation in schizophrenia.

The largest published schizophrenia GWAS, that from the Schizophrenia Working Group of the Psychiatric Genomics Consortium (PGC), identified 108 genome-wide significant (GWS) loci and unequivocally demonstrated the value of increasing sample sizes for discovery in schizophrenia genetics research⁵. Here, we report a large, phenotypically homogeneous GWA

study of schizophrenia which, when combined with previous published data, identifies novel facets of genetic architecture and biology, and demonstrates that the evolutionary process of background selection can explain the persistence of common risk alleles in the population.

RESULTS

GWAS and Meta-analysis

We obtained genome-wide genotype information for schizophrenia cases from the UK (the CLOZUK sample), which we combined with control datasets obtained from public repositories or through collaboration. The final sample size was of 11,260 cases and 24,542 controls (5,220 cases and 18,823 controls not in previous schizophrenia GWAS; **Methods; Supplementary Figure 1; Supplementary Figure 2**). At a genome wide level, the association statistics indicated that the common variant architecture in the CLOZUK sample was highly correlated with an independent sample of 29,415 cases and 40,101 controls from the PGC (genetic correlation = 0.954 ± 0.030 ; $p = 6.63 \times 10^{-227}$) and this was further confirmed by polygenic risk score and trend test analyses across the datasets at a range of association p-value thresholds (**Methods and Supplementary Table 1 and 2**).

Meta-analysis of the CLOZUK and the independent PGC dataset, excluding related and overlapping samples (total 40,675 cases and 64,643 controls; **Supplementary Figure 1**), identified 179 independent GWS SNPs ($p < 5 \times 10^{-8}$, **Supplementary Table 3**) mapping to 145 independent loci (**Methods, Figure 1, Supplementary Table 4**). The 145 associated loci include 93 of those that were GWS in the study of the PGC, the majority of which showed a strengthened association (**Supplementary Figure 4, Supplementary Table 5**). This does not imply the remaining 15 PGC loci were false positives, rather this reflects the expected inflation of effect sizes for GWS SNPs in incompletely powered studies, and as we

demonstrate, is consistent with all 108 PGC loci representing true positives (see **Supplementary Note**). Of the 52 loci not identified by the PGC, two have been reported as genome-wide significant in other studies: the locus at ZEB2¹⁴ and a locus on chromosome 8 (38.0-38.3 MB)¹⁵.

In further independent samples (5,662 cases and 154,224 controls); 43 of the 50 GWS index SNPs showed the same pattern of allelic association, a level that far surpasses chance ($p=1.05 \times 10^{-7}$). Despite the modest number of cases in these samples, 18 of the 50 index alleles reached nominal significance ($p < 0.05$), again implausible by chance ($p=1.46 \times 10^{-11}$). None demonstrated evidence for heterogeneity of effect (**Methods, Supplementary Table 6**).

Mutation intolerant genes

Recent studies have shown that mutation intolerant genes capture much of the rare variant architecture of neurodevelopmental disorders such as autism, intellectual disability and developmental delay as well as schizophrenia¹⁶⁻¹⁹. Here, we show that for schizophrenia, this also holds for common variation. Using gene set analysis in MAGMA²⁰, loss-of-function (LoF) intolerant genes (N=3,230) as defined by the Exome Aggregation Consortium (ExAC)²¹ using their gene-level constraint metric $pLI \geq 0.9$, were enriched for schizophrenia common variant associations in comparison with all other annotated genes ($p=4.1 \times 10^{-16}$).

It has been shown that pLI is correlated with gene expression across tissues, including brain²¹, which raises the possibility that the LoF-intolerant gene enrichment in schizophrenia may reflect enrichment for signal in genes expressed in the brain. However, LoF-intolerant gene set enrichment was robust to the inclusion of “brain expressed” (N=10,360) and “brain specific” (N=2,647) gene sets¹⁹ as covariates in the analysis ($p=1.89 \times 10^{-10}$) or to controlling for FPKM gene expression values ($p=1.03 \times 10^{-14}$) in brain²².

It has been suggested that clustering of risk alleles in mutation intolerant genes is a hallmark of early-onset traits under natural selection^{23,24}. However, LoF-intolerant genes are known to be enriched for SNPs identified as genome-wide significant in GWAS studies (as listed in the NHGRI-EBI GWAS Catalog²⁵) and for broad categories of disorders²¹. To examine whether our finding is a property of polygenic disorders in general, we obtained summary genetic data from a neuropsychiatric and non-psychiatric late-onset disorder (Alzheimer's disease, type-2 diabetes) and a psychological trait (Neuroticism), each of which has been shown to be under minimal selective pressure (see **Methods**). These other phenotypes show at best a weak signal for enrichment of the LoF-intolerant gene set in the MAGMA analysis, not comparable to that seen in schizophrenia (Alzheimer's disease $p=0.008$, type-2 diabetes $p=0.016$, Neuroticism $p=0.066$).

To quantify the contribution of SNPs within LoF-intolerant genes to schizophrenia SNP-based heritability (h^2_{SNP}) we used partitioned LDSC²⁶ (**Supplementary Table 7**). Overall, genic SNPs account for 64% of h^2_{SNP} , a 1.23-fold enrichment proportional to their SNP content ($p=5.93 \times 10^{-14}$). Consistent with the analysis using MAGMA, h^2_{SNP} was enriched in LoF-intolerant genes (2.01-fold; $p=2.78 \times 10^{-24}$), which explained 30% of all h^2_{SNP} (equating to 47% of all genic h^2_{SNP}). In contrast, genes classed as non LoF-intolerant ($pLI < 0.9$) were significantly depleted for h^2_{SNP} relative to their SNP content (0.90-fold; $p=5.86 \times 10^{-3}$), although in absolute terms, SNPs in these genes accounted for 34% of h^2_{SNP} . A finer scale analysis of the relationship between LoF intolerance scores and enrichment for association showed that enrichment is restricted to genes with a pLI score above 0.9, precisely those defined as “LoF-intolerant” (**Supplementary Figure 5**).

Common risk alleles in regions under background selection

Our finding that LoF-intolerant genes are enriched for common risk variants raises the question of how such alleles are found at common frequencies in the population. While the

contribution of ultra-rare variation in functionally important genes to disorders associated with low fecundity can be accounted for by *de novo* mutation^{16,19,27}, this cannot explain the persistence of common alleles. To address this question, we used partitioned LDSR to test the relationship between schizophrenia associated alleles and SNP-based signatures of natural selection. These included measures of positive selection, background selection, and Neanderthal introgression. We examined the heritability of SNPs after thresholding them at extreme values for these metrics (top 2%, 1% and 0.5%), including in the baseline model annotation sets such as LoF-intolerant genes and genomic regions with extreme linkage disequilibrium patterns (**Methods**).

We observed strong evidence for schizophrenia h^2_{SNP} enrichment in SNPs under strong background selection (BGS), which was consistent across all the thresholds we examined (**Table 1**). We also found a significant depletion of h^2_{SNP} in SNPs subject to positive selection as indexed by the CLR statistic. These two results are mutually consistent, as the calculation of the CLR statistic explicitly controls for the effect of BGS²⁸. This suggests that SNPs under positive selection, but under weak or no BGS, are depleted for association with schizophrenia. No significant relationship between h^2_{SNP} and other positive selection or Neanderthal introgression measures was found after correction for multiple testing (**Table 1**). An LDSR analysis treating BGS measures as a quantitative trait, rather than as a binary one, confirmed that the relationship between BGS and schizophrenia association was not due to the imposition of arbitrary thresholds to define strong BGS ($p=7.73 \times 10^{-11}$). We also note that the τ_c statistic of the LDSC model is significant for BGS, in both binary ($p=0.041$) and quantitative ($p=0.023$) analyses (**Supplementary Table 8**). The τ_c statistic indicates the enrichment of BGS after controlling for all other annotations in the model (including LoF-intolerant genes)²⁶, and thus represents a robust and conservative test for the BGS enrichment.

The above analyses accounts for a possible confounding relationship between LoF intolerance and BGS. To illustrate this more clearly, we binned the BGS intensities into four categories of increasing score, and classified SNPs in these bins according to whether they are in LoF-intolerant genes, “all other” genes sets and a non-genic set (**Supplementary Figure 6**). Note that the lower boundary of the top bin (BGS intensity > 0.75) corresponds approximately to the top 2% BGS threshold in **Table 1** and is equivalent to a reduction in effective population size estimated at each SNP of 75% or more²⁹. We found significant heritability enrichment across all BGS intensity intervals in LoF-intolerant genes that increased progressively with higher intensity scores. Importantly, we also found heritability enrichment for SNPs under BGS pressure in genes that are not LoF intolerant, restricted to the highest BGS intensity bin. Indeed the highest BGS intensity bin in non-LoF genes was enriched for heritability at a level roughly equivalent to all LoF genes. These findings point to BGS and LoF intolerance as making at least partially independent contributions to heritability enrichment in schizophrenia. In contrast, none of the phenotypes we selected on the basis of their minimal impact on fecundity (Alzheimer’s disease, type-2 diabetes, neuroticism) showed significant BGS enrichment for heritability using either the BGS τ_c statistic of the LDSC model (minimum $p > 0.24$), or when specifically testing regions of high BGS intensity in genes that are tolerant ($pLI < 0.9$) of functional mutations (minimum $p > 0.40$).

Systems genomics

Using MAGMA, we undertook a primary analysis of 134 central nervous system related gene sets we have previously shown captures the excess CNV burden in schizophrenia³⁰. In a GWAS context, we now show that collectively, this group of gene sets captures a disproportionately high fraction of h^2_{SNP} (30% of total heritability; enrichment = 1.63; $p = 8.57 \times 10^{-13}$; 46% of genic heritability; **Supplementary Table 8**). Of the 134 sets, 54 were nominally significant of which 12 survived multiple-testing correction (family-wise error rate

(FWER) $p < 0.05$, **Supplementary Table 9**), with no notable association for gene sets such as the ARC protein complex and NMDAR protein network, that we have previously implicated in rare variant studies^{30,31}. Stepwise conditional analysis, adjusting sequentially for the more strongly associated gene sets, resulted in six gene sets that were independently associated with schizophrenia (**Table 2** and **Data Supplement**). These extend from low-level molecular and sub-cellular processes to broad behavioural phenotypes. The most strongly associated gene set is constituted by the targets of the Fragile X Mental Retardation Protein (FMRP)³². FMRP is a neuronal RNA-binding protein that interacts with polyribosomal mRNAs (the 842 target transcripts of this gene set³²) and is thought to act by inhibiting translation of target mRNAs, including many transcripts of pre- and post-synaptic proteins. The FMRP target set has been shown to be enriched for rare mutational burden in *de novo* exome sequencing studies of autism³³ and intellectual disability³¹. In schizophrenia studies, it has also been shown to be nominally significantly enriched for association signal in sequencing studies^{8,31} and in GWAS^{5,8} but only inconsistently in studies of copy number variation^{30,34}. Here we provide the strongest evidence to date for the enrichment of this gene set in schizophrenia.

We highlight another five gene sets that are independently associated with schizophrenia. Three of these derive from the Mouse Genome Informatics database³⁵ and relate to behavioural and neurophysiological correlates of learning; Abnormal Behaviour (MP:0004924), Abnormal Nervous System Electrophysiology (MP:0002272) and Abnormal Long Term Potentiation (MP:0002207). We note that two of these gene sets (MP:0004924 and MP:0002207) were among the five most enriched of 134 gene sets tested in a recent schizophrenia CNV analysis³⁰. The remaining two independently associated genes sets were voltage-gated calcium channel complexes³⁶ and the 5-HT_{2C} receptor complex³⁷. The calcium channel finding confirms extensive evidence from common and rare variant studies

implicating calcium channel genes in schizophrenia^{5,8}, including a novel GWAS locus in *CACNA1D* identified in our meta-analysis. Whilst there is less convergent evidence in support of the involvement of the 5-HT_{2C} receptor complex in schizophrenia, the fact that we identify independent association for this gene set implicates these genes in schizophrenia pathophysiology and potentially rejuvenates a previous avenue of 5-HT_{2C} ligand therapeutic endeavour in schizophrenia research³⁸. However we interpret this result with caution given the small size of this gene set and the fact that a number of its genes encode synaptic proteins that are structurally related to other receptor complexes³⁷, not only 5-HT_{2C}.

Systems genomics and mutation intolerant genes

The LoF-intolerant genes and the six conditionally independent (“significant”) CNS-related gene sets together account for 39% of schizophrenia SNP-based heritability ($p=5.07 \times 10^{-26}$), equating to 61% of genic heritability (**Figure 2A; Supplementary Table 7**). This is likely to be an underestimation of the true effect of these gene sets since distal non-genic regulatory elements (not included in this analysis) will add to the heritability explained by these genes. In examining the relationship between the LoF-intolerant and CNS-related gene sets (**Figure 2A**), genes belonging to both categories were the most highly enriched (2.6-fold; $p=7.90 \times 10^{-15}$), although LoF-intolerant genes that were not annotated to our significant CNS gene sets still displayed enrichment for SNP-based heritability (1.74-fold; $p=9.77 \times 10^{-10}$), while genes that were in the significant CNS gene sets but had $pLI < 0.9$ showed more modest enrichment (1.39-fold; $p=6.05 \times 10^{-4}$). Notably genes outside these categories were depleted in heritability relative to their SNP content (enrichment=0.79, $p=1.82 \times 10^{-7}$).

This general pattern remained when we focussed on the six significant CNS gene sets individually, in that the enrichment in these gene sets derives primarily from their intersection with LoF-intolerant genes (**Figure 2B**). Indeed, only the targets of FMRP showed significant enrichment for SNPs in genes that are not LoF intolerant (2.06-fold; $p=4.23 \times 10^{-5}$).

Data-driven gene set analysis

To set the systems genomics results in context, and to ensure we were not missing enrichment in other gene sets by our hypothesis driven approach, we undertook a purely data-driven analysis of a larger comprehensive annotation of gene sets from multiple public databases, totalling 6,677 gene sets (**Methods, Supplementary Table 10**). Six gene sets survived FWER correction for the full 6,677 gene sets and showed independence through conditional analyses. The LoF-intolerant gene set was the most strongly enriched followed by the two most strongly associated functional gene sets we had specified in our hypothesis-driven CNS gene set analysis (FMRP targets and MGI Abnormal Behaviour genes). The other three sets were calcium ion import (GO:0070509), membrane depolarisation during action potential (GO:0086010) and synaptic transmission (GO:0007268). These are highly overlapping with the independently associated sets from our primary CNS systems genomic analysis. Indeed if we repeat the data-driven comprehensive gene set analysis whilst adjusting for the six independently associated CNS gene sets, then the only surviving enrichment term is the LoF-intolerant genes. These results are consistent with those from CNV analysis³⁰ in that they do not support annotations other than those related to CNS function, and demonstrate that hypothesis based analysis to maximise power does not substantially impact on the overall pattern of results.

Identifying likely candidates within associated loci

To identify SNPs and genes which might be causally linked to the GWS associations, we used FINEMAP³⁹ to identify credibly causal alleles (those with a cumulative posterior probability for a locus of at least 95%) and functionally annotated these alleles using ANNOVAR⁴⁰. This identified 6,105 credible SNPs across 144 GWS loci, excluding the MHC region (**Methods, Supplementary Table 11**). From these, we defined a highly credible set of SNPs (N=25) as those that are more likely to explain the associations than all other

SNPs combined (i.e. with a FINEMAP posterior probability greater than 0.5). Of these, 14 mapped to genes based on putative functionality (exonic SNPs that cause non-synonymous or splice variations or promoter SNPs, n=6) or mapped to regions identified as likely regulatory elements (n=8) through chromosome conformation analysis performed in tissue from the developing brain using Hi-C⁴¹ physical interactions (**Methods; Supplementary Table 12**). One of the implicated alleles is a nonsynonymous variant in the manganese and zinc transporter gene *SLC39A8*. Nonsynonymous variants in this gene have been associated with severe neurodevelopmental disorders and deficiencies of SLC39A8 with related impaired glycosylation⁴², highlighting a mechanism of therapeutic potential.

We also applied Summary-data based Mendelian Randomisation (SMR) analysis⁴³ to the data in concert with the dorsolateral prefrontal cortex eQTL data from the CommonMind Consortium⁴⁴ aiming to identify variants that might be causally linked through expression changes of specific genes. (**Methods, Supplementary Table 13**). After applying a conservative threshold ($p_{\text{HEIDI}} > 0.05$) which prioritises those co-localised signals due to a single causal variant⁴³, we identified 22 candidates at 19 loci with a false discovery rate $p < 0.05$.

In total, the combination of FINEMAP, Hi-C and SMR analyses assigned potentially causal genes at 33 GWS loci and implicated a single gene at 27 of these loci. However, the analyses intersect for a single gene, ZNF823, indicating the need for more comprehensive functional genomic annotations in CNS relevant tissues.

DISCUSSION

In the largest genetic study of schizophrenia to date, we explore the genomic architecture of, and the evolutionary pressures on, common variants associated with the disorder. Our study

provides the first evidence linking common variation in LoF-intolerant genes to risk of developing schizophrenia and demonstrates that these genes account for a substantial proportion (30%) of schizophrenia SNP-based heritability. Systems genomic analysis highlights six gene sets that are independently associated with schizophrenia, and point to molecular, physiological and behavioural pathways involved in schizophrenia pathogenesis.

Given mutation intolerance is due to high selection pressure^{21,23,24}, our finding that schizophrenia risk variants that persist at common allele frequencies are enriched in loss-of-function intolerant genes might appear counter-intuitive. However, novel evidence presented here suggests this can be reconciled by background selection (BGS) which is a consequence of purifying selection in regions of low recombination^{45,46}. In such regions, recurrent selection against deleterious variants causes haplotypes to be removed from the gene pool, which reduces genetic diversity in a manner equivalent to a reduction in effective population size⁴⁷. This in turn impairs the efficiency of the selection process, allowing alleles with small deleterious effects to rise in frequency by drift⁴⁸. Such a consequence of purifying selection has been shown to be compatible with the genomic architecture of complex human traits⁴⁹ and to influence phenotypes in model organisms⁵⁰. We have explicitly modelled this effect (both theoretically and via simulations; **Supplementary Note**) and provide strong evidence for the feasibility of this effect as explanatory for the effect sizes seen for common alleles in schizophrenia.

We did not find enrichment for any measure of positive selection or Neanderthal introgression. A recent study explained a negative correlation between schizophrenia associations and metrics indicative of a Neanderthal selective sweep as evidence for positive selection or polygenic adaptation in schizophrenia¹². We do not find any significant correlation in our model, which addresses the contribution of BGS, and hence our results are not consistent with large contributions of positive selection to the genetic architecture of

schizophrenia (**Table 1**). Indeed positive selection is not widespread in humans, as reported by other studies that explicitly considered or accounted for BGS^{28,51}. Polygenic adaptation, the co-occurrence of many subtle allele frequency shifts at loci influencing complex traits⁵², remains an intriguing possibility, but has not been implicated in psychiatric phenotypes, including schizophrenia, in recent analyses^{53,54}. In contrast, BGS has been proposed as a mechanism driving Human-Neanderthal incompatibilities, as regions with stronger estimated BGS have lower estimated Neanderthal introgression⁵⁵. We therefore conclude that the bulk of the BGS signal we obtain is unlikely to be influenced by positive selection²⁹, challenging theories of selective advantage of schizophrenia risk alleles to explain their high population frequencies.

AUTHOR CONTRIBUTIONS

A.F.P. curated and processed genetic data, performed statistical analyses, contributed to the interpretation of results and participated in the primary drafting of the manuscript.

P.H., A.J.P., V.E-P., A.C. and E.S. performed statistical analyses, contributed to the interpretation of results and participated in the primary drafting of the manuscript.

S.R. curated and processed genetic data and participated in the primary drafting of the manuscript.

N.C. and M.L.H. contributed to the interpretation of results and participated in the primary drafting of the manuscript.

S.E.L., S.B. and A.L. participated in the recruitment of participants for the study and curated and managed their phenotypic information.

D.C., J.H, L.H, E.R. and G.K. contributed and curated data used in the statistical analyses.

K.M. managed the laboratory and genotyping procedures in Cardiff University.

J. H. M, D. A. C. and D.R. supervised the recruitment of the participants for the study.

S. A. M. managed the genotyping of samples for the study.

N. R. W. contributed genotypes of control samples and participated in the primary drafting of the manuscript.

D.H. G, L. M. H., D. M. R., P. S., E. A. S. and H.W. performed statistical analyses and contributed to the interpretation of results.

M. J. O. and M. C. O'D. conceived and supervised the project, contributed to the interpretation of results and participated in the primary drafting of the manuscript.

J. T. R. W. conceived and supervised the project, led the recruitment of the participants and sample acquisition for the study, performed statistical analysis, contributed to the interpretation of results and participated in the primary drafting of the manuscript.

All other authors contributed genotypes of control samples or summary statistics of replication samples.

All authors had the opportunity to review and comment on the manuscript and all approved the final manuscript.

COMPETING FINANCIAL INTERESTS

D. A. C. is a full-time employee and stockholder of Eli Lilly and Company. The remaining authors declare no conflicts of interest.

REFERENCES

1. Owen, M.J., Sawa, A. & Mortensen, P.B. Schizophrenia. *Lancet* (2016).
2. Thornicroft, G. Physical health disparities and mental illness: the scandal of premature mortality. *The British Journal of Psychiatry* **199**, 441-442 (2011).
3. Olfson, M., Gerhard, T., Huang, C., Crystal, S. & Stroup, T. Premature mortality among adults with schizophrenia in the united states. *JAMA Psychiatry* **72**, 1172-1181 (2015).
4. Morgan, C. *et al.* Reappraising the long-term course and outcome of psychotic disorders: the AESOP-10 study. *Psychological Medicine* **44**, 2713-2726 (2014).
5. Schizophrenia Working Group of the Psychiatric Genomics Consortium. Biological insights from 108 schizophrenia-associated genetic loci. *Nature* **511**, 421-427 (2014).
6. Singh, T. *et al.* Rare loss-of-function variants in SETD1A are associated with schizophrenia and developmental disorders. *Nature Neuroscience* **19**, 571-7 (2016).

7. Rees, E. *et al.* Analysis of copy number variations at 15 schizophrenia-associated loci. *Br J Psychiatry* **204**, 108-14 (2014).
8. Purcell, S.M. *et al.* A polygenic burden of rare disruptive mutations in schizophrenia. *Nature* **506**, 185-190 (2014).
9. Power, R.A. *et al.* Fecundity of patients with schizophrenia, autism, bipolar disorder, depression, anorexia nervosa, or substance abuse vs their unaffected siblings. *JAMA psychiatry* **70**, 22-30 (2013).
10. Huxley, J., Mayr, E., Osmond, H. & Hoffer, A. Schizophrenia as a Genetic Morphism. *Nature* **204**, 220-221 (1964).
11. Shaner, A., Miller, G. & Mintz, J. Schizophrenia as one extreme of a sexually selected fitness indicator. *Schizophrenia Research* **70**, 101-109 (2004).
12. Srinivasan, S. *et al.* Genetic Markers of Human Evolution Are Enriched in Schizophrenia. *Biological Psychiatry* **80**, 284–292 (2016).
13. Uher, R. The role of genetic variation in the causation of mental illness: an evolution-informed framework. *Mol Psychiatry* **14**, 1072-1082 (2009).
14. Ripke, S. *et al.* Genome-wide association analysis identifies 13 new risk loci for schizophrenia. *Nature Genetics* **45**, 1150-1159 (2013).
15. Shi, Y. *et al.* Common variants on 8p12 and 1q24.2 confer risk of schizophrenia. *Nature Genetics* **43**, 1224-1227 (2011).
16. McRae, J.F. *et al.* Prevalence and architecture of de novo mutations in developmental disorders. *Nature* **542**, 433–438 (2017).
17. Kosmicki, J.A. *et al.* Refining the role of de novo protein-truncating variants in neurodevelopmental disorders by using population reference samples. *Nature Genetics* **49**, 504–510 (2017).
18. Samocha, K.E. *et al.* A framework for the interpretation of de novo mutation in human disease. *Nature Genetics* **46**, 944-950 (2014).
19. Genovese, G. *et al.* Increased burden of ultra-rare protein-altering variants among 4,877 individuals with schizophrenia. *Nature Neuroscience* **19**, 1433-1441 (2016).
20. de Leeuw, C.A., Mooij, J.M., Heskes, T. & Posthuma, D. MAGMA: Generalized gene-set analysis of GWAS data. *PLoS Comput Biol* **11**, e1004219 (2015).
21. Lek, M. *et al.* Analysis of protein-coding genetic variation in 60,706 humans. *Nature* **536**, 285–291 (2016).
22. Fagerberg, L. *et al.* Analysis of the Human Tissue-specific Expression by Genome-wide Integration of Transcriptomics and Antibody-based Proteomics. *Molecular & Cellular Proteomics* **13**, 397-406 (2014).

23. Smith, N.G.C. & Eyre-Walker, A. Human disease genes: patterns and predictions. *Gene* **318**, 169-175 (2003).
24. Blekhman, R. *et al.* Natural selection on genes that underlie human disease susceptibility. *Current biology : CB* **18**, 883-889 (2008).
25. Welter, D. *et al.* The NHGRI GWAS Catalog, a curated resource of SNP-trait associations. *Nucleic acids research* **42**, D1001-D1006 (2014).
26. Finucane, H.K. *et al.* Partitioning heritability by functional annotation using genome-wide association summary statistics. *Nature Genetics* **47**, 1228-1235 (2015).
27. Takata, A., Ionita-Laza, I., Gogos, Joseph A., Xu, B. & Karayiorgou, M. De Novo Synonymous Mutations in Regulatory Elements Contribute to the Genetic Etiology of Autism and Schizophrenia. *Neuron* **89**, 940-947 (2016).
28. Huber, C.D., DeGiorgio, M., Hellmann, I. & Nielsen, R. Detecting recent selective sweeps while controlling for mutation rate and background selection. *Molecular Ecology* **25**, 142-156 (2016).
29. McVicker, G., Gordon, D., Davis, C. & Green, P. Widespread Genomic Signatures of Natural Selection in Hominid Evolution. *PLoS Genet* **5**, e1000471 (2009).
30. Pocklington, A.J. *et al.* Novel Findings from CNVs Implicate Inhibitory and Excitatory Signaling Complexes in Schizophrenia. *Neuron* **86**, 1203-14 (2015).
31. Fromer, M. *et al.* De novo mutations in schizophrenia implicate synaptic networks. *Nature* **506**, 179-184 (2014).
32. Darnell, J.C. *et al.* FMRP stalls ribosomal translocation on mRNAs linked to synaptic function and autism. *Cell* **146**, 247-61 (2011).
33. Iossifov, I. *et al.* De Novo Gene Disruptions in Children on the Autistic Spectrum. *Neuron* **74**, 285-299 (2012).
34. Szatkiewicz, J.P. *et al.* Copy number variation in schizophrenia in Sweden. *Molecular Psychiatry* **19**, 762-73 (2014).
35. Blake, J.A., Bult, C.J., Eppig, J.T., Kadin, J.A. & Richardson, J.E. The Mouse Genome Database: integration of and access to knowledge about the laboratory mouse. *Nucleic Acids Research* **42**, D810-7 (2014).
36. Müller, C.S. *et al.* Quantitative proteomics of the Cav2 channel nano-environments in the mammalian brain. *Proceedings of the National Academy of Sciences* **107**, 14950-14957 (2010).
37. Becamel, C. *et al.* Synaptic multiprotein complexes associated with 5-HT(2C) receptors: a proteomic approach. *The EMBO journal* **21**, 2332-42 (2002).
38. Liu, J. *et al.* Prediction of Efficacy of Vabicaserin, a 5-HT(2C) Agonist, for the Treatment of Schizophrenia Using a Quantitative Systems Pharmacology Model. *CPT: Pharmacometrics & Systems Pharmacology* **3**, e111 (2014).

39. Benner, C. *et al.* FINEMAP: efficient variable selection using summary data from genome-wide association studies. *Bioinformatics* **32**, 1493-1501 (2016).
40. Wang, K., Li, M. & Hakonarson, H. ANNOVAR: functional annotation of genetic variants from high-throughput sequencing data. *Nucleic acids research* **38**, e164-e164 (2010).
41. Won, H. *et al.* Chromosome conformation elucidates regulatory relationships in developing human brain. *Nature* **538**, 523-527 (2016).
42. Park, J.H. *et al.* SLC39A8 Deficiency: A Disorder of Manganese Transport and Glycosylation. *American Journal of Human Genetics* **97**, 894-903 (2015).
43. Zhu, Z. *et al.* Integration of summary data from GWAS and eQTL studies predicts complex trait gene targets. *Nature Genetics* **48**, 481-487 (2016).
44. Fromer, M. *et al.* Gene expression elucidates functional impact of polygenic risk for schizophrenia. *Nat Neurosci* **19**, 1442-1453 (2016).
45. Charlesworth, B. The Effects of Deleterious Mutations on Evolution at Linked Sites. *Genetics* **190**, 5-22 (2012).
46. Charlesworth, B., Betancourt, A.J., Kaiser, V.B. & Gordo, I. Genetic Recombination and Molecular Evolution. *Cold Spring Harbor Symposia on Quantitative Biology* **74**, 177-186 (2009).
47. Comeron, J.M., Williford, A. & Kliman, R.M. The Hill-Robertson effect: evolutionary consequences of weak selection and linkage in finite populations. *Heredity* **100**, 19-31 (2007).
48. Charlesworth, B. Background Selection 20 Years on: The Wilhelmine E. Key 2012 Invitational Lecture. *Journal of Heredity* (2013).
49. North, T.L. & Beaumont, M.A. Complex trait architecture: the pleiotropic model revisited. *Scientific Reports* **5**, 9351 (2015).
50. Rockman, M.V., Skrovanek, S.S. & Kruglyak, L. Selection at linked sites shapes heritable phenotypic variation in *C. elegans*. *Science* **330**, 372-6 (2010).
51. Vitti, J.J., Grossman, S.R. & Sabeti, P.C. Detecting Natural Selection in Genomic Data. *Annual Review of Genetics* **47**, 97-120 (2013).
52. Stephan, W. Signatures of positive selection: from selective sweeps at individual loci to subtle allele frequency changes in polygenic adaptation. *Molecular ecology* **25**, 79-88 (2016).
53. Field, Y. *et al.* Detection of human adaptation during the past 2,000 years. *Science* (2016).
54. Key, F.M., Fu, Q., Romagne, F., Lachmann, M. & Andres, A.M. Human adaptation and population differentiation in the light of ancient genomes. *Nat Commun* **7**(2016).

55. Harris, K. & Nielsen, R. The Genetic Cost of Neanderthal Introgression. *Genetics* **203**, 881-891 (2016).

ACKNOWLEDGEMENTS

General

This project has received funding from the European Union's Seventh Framework Programme for research, technological development and demonstration under grant agreement n° 279227 (CRESTAR Consortium). The work at Cardiff University was funded by Medical Research Council (MRC) Centre (MR/L010305/1), Program Grant (G0800509) and the European Community's Seventh Framework Programme HEALTH-F2-2010-241909 (Project EU-GEI). U.D. received funding by the German Research Foundation (DFG, grant FOR2107 DA1151/5-1; SFB-TRR58, Project C09) and the Interdisciplinary Center for Clinical Research (IZKF) of the medical faculty of Münster (grant Dan3/012/17). E.S. and A.C. received funding from Agencia Estatal de Investigación (AEI; CGL2016-75904-C2-1-P), Xunta de Galicia (ED431C 2016-037) and Fondo Europeo de Desarrollo Regional (FEDER). The iPSYCH and GEMS2 teams acknowledge funding from The Lundbeck Foundation (grant no R102-A9118 and R155-2014-1724), the Stanley Medical Research Institute, an Advanced Grant from the European Research Council (project no: 294838), the Danish Strategic Research Council and grants from Aarhus University to the iSEQ and CIRRAU centers.

Case data

We thank the participants and clinicians who took part in the CardiffCOGS study. For the CLOZUK2 sample we thank Leyden Delta for supporting the sample collection, anonymisation and data preparation (particularly Marinka Helthuis, John Jansen, Karel Jollie and Anouschka Colson), Magna Laboratories, UK (Andy Walker) and, for CLOZUK1,

Novartis and The Doctor's Laboratory staff for their guidance and cooperation. We acknowledge Lesley Bates, Catherine Bresner and Lucinda Hopkins, at Cardiff University, for laboratory sample management. We acknowledge Wayne Lawrence and Mark Einon, at Cardiff University, for support with the use and setup of computational infrastructures.

Control data

A full list of the investigators who contributed to the generation of the Wellcome Trust Case Control Consortium (WTCCC) data is available from its URL. Funding for the project was provided by the Wellcome Trust (WT) under award 076113. The UK10K project was funded by the Wellcome Trust award WT091310. Venous blood collection for the 1958 Birth Cohort (NCDS) was funded by the UK's Medical Research Council (MRC) grant G0000934, peripheral blood lymphocyte preparation by Juvenile Diabetes Research Foundation (JDRF) and WT and the cell-line production, DNA extraction and processing by WT grant 06854/Z/02/Z. Genotyping was supported by WT (083270) and the European Union (EU; ENGAGE: HEALTH-F4-2007- 201413). The UK Blood Services Common Controls (UKBS-CC collection) was funded by WT (076113/C/04/Z) and by the National Institute for Health Research (NIHR) programme grant to NHS Blood and Transplant authority (NHSBT; RP-PG-0310-1002). NHSBT also made possible the recruitment of the Cardiff Controls, from participants who provided informed consent. Generation Scotland (GS) received core funding from the Chief Scientist Office of the Scottish Government Health Directorates CZD/16/6 and the Scottish Funding Council HR03006. Genotyping of the GS:SFHS samples was carried out by the Genetics Core Laboratory at the WT Clinical Research Facility, Edinburgh, Scotland and was funded by the MRC. The Type 1 Diabetes Genetics Consortium (T1DGC; EGA dataset EGAS00000000038) is a collaborative clinical study sponsored by the National Institute of Diabetes and Digestive and Kidney Diseases (NIDDK), National Institute of Allergy and Infectious Diseases (NIAID), National Human Genome Research Institute

(NHGRI), National Institute of Child Health and Human Development (NICHD), and JDRF. The People of the British Isles project (POBI) is supported by WT (088262/Z/09/Z). TwinsUK is funded by WT, MRC, EU, NIHR-funded BioResource, Clinical Research Facility and Biomedical Research Centre based at Guy's and St Thomas' NHS Foundation Trust in partnership with King's College London. Funding for the QIMR samples was provided by the Australian National Health and Medical Research Council (241944, 339462, 389875, 389891, 389892, 389927, 389938, 442915, 442981, 496675, 496739, 552485, 552498, 613602, 613608, 613674, 619667), the Australian Research Council (FT0991360, FT0991022), the FP-5 GenomeUtwinn Project (QLG2-CT- 2002-01254) and the US National Institutes of Health (NIH; AA07535, AA10248, AA13320, AA13321, AA13326, AA14041, MH66206, DA12854, DA019951), and the Center for Inherited Disease Research (Baltimore, MD, USA). TEDS is supported by a program grant from the MRC (G0901245-G0500079), with additional support from the NIH (HD044454; HD059215). In the GERAD1 Consortium, Cardiff University was supported by WT, MRC, Alzheimer's Research UK (ARUK) and the Welsh Government. King's College London acknowledges support from the MRC. The University of Belfast acknowledges support from ARUK, Alzheimer's Society, Ulster Garden Villages, N.Ireland R&D Office and the Royal College of Physicians/Dunhill Medical Trust. Washington University was funded by NIH grants, Barnes Jewish Foundation and the Charles and Joanne Knight Alzheimer's Research Initiative. The Bonn group was supported by the German Federal Ministry of Education and Research (BMBF), Competence Network Dementia and Competence Network Degenerative Dementia, and by the Alfred Krupp von Bohlen und Halbach-Stiftung.

CRESTAR Consortium Members

Evanthia Achilla¹, Esben Agerbo^{2,3}, Cathy L. Barr⁴, Theresa Wimberly Böttger², Gerome Breen^{5,6}, Dan Cohen⁷, David A. Collier^{5,8}, Sarah Curran^{9,10}, Emma Dempster¹¹, Danai Dima⁵,

Ramon Sabes-Figuera¹, Robert J. Flanagan¹², Sophia Frangou¹³, Josef Frank¹⁴, Christiane Gasse^{2,3}, Fiona Gaughran¹⁵, Ina Giegling¹⁶, Jakob Grove^{18,19}, Eilis Hannon¹¹, Annette M. Hartmann¹⁶, Barbara Hei  er¹⁹, Marinka Helthuis²⁰, Henriette Thisted Horsdal², Oddur Ingimarsson²¹, Karel Jollie²⁰, James L. Kennedy²², Ole K  hler²³, Bettina Konte¹⁶, Maren Lang¹⁴, Sophie Legge²⁴, Cathryn Lewis⁵, James MacCabe¹⁵, Anil K. Malhotra²⁵, Paul McCrone¹, Sandra M. Meier², Jonathan Mill^{5,11}, Ole Mors²³, Preben Bo Mortensen², Markus M. N  then²⁶, Michael C. O'Donovan²⁴, Michael J. Owen²⁴, Antonio F. Pardi  as²⁴, Carsten B. Pedersen^{2,3}, Marcella Rietschel¹⁴, Dan Rujescu¹⁶, Ameli Schwalber¹⁹, Engilbert Sigurdsson²¹, Holger J. S  rensen²⁷, Benjamin Spencer²⁸, Hreinn Stefansson²⁹, Henrik St  vring³, Jana Strohmaier¹⁴, Patrick Sullivan^{30,31}, Evangelos Vassos⁵, Moira Verbelen⁵, James T. R. Walters²⁴, Thomas Werge²⁷.

¹Centre for Economics of Mental and Physical Health, Health Service and Population Research Department, Institute of Psychiatry, King's College London, London, UK.

²National Centre for Register-Based Research, Department of Economics and Business, School of Business and Social Sciences, Aarhus University, Aarhus, Denmark. ³Centre for Integrated Register-based Research, CIRRAU, Aarhus University, Aarhus, Denmark.

⁴Toronto Western Research Institute, University Health Network, Toronto, Canada. ⁵MRC Social, Genetic and Developmental Psychiatry Centre, Institute of Psychiatry, Psychology and Neuroscience, King's College London, London, UK. ⁶NIHR Biomedical Research Centre for Mental Health, Maudsley Hospital and Institute of Psychiatry, Psychology and Neuroscience, King's College London, London, UK. ⁷Department of Community Mental Health, Mental Health Organization North-Holland North, Heerhugowaard, The Netherlands.

⁸Discovery Neuroscience Research, Eli Lilly and Company Ltd, Lilly Research Laboratories, Erl Wood Manor, Surrey, UK. ⁹Department of Child & Adolescent Psychiatry, Institute of Psychiatry, Psychology & Neuroscience, King's College London, London, UK. ¹⁰Brighton

and Sussex Medical School, University of Sussex, Brighton, UK. ¹¹University of Exeter Medical School, RILD, University of Exeter, Exeter, UK. ¹²Toxicology Unit, Department of Clinical Biochemistry, King's College Hospital NHS Foundation Trust, London, UK. ¹³Clinical Neurosciences Studies Center, Department of Psychiatry, Icahn School of Medicine at Mount Sinai, New York, USA. ¹⁴Department of Genetic Epidemiology in Psychiatry, Central Institute of Mental Health, Medical Faculty Mannheim/Heidelberg University, Mannheim, Germany. ¹⁵Department of Psychosis Studies, Institute of Psychiatry, Psychology and Neuroscience, King's College London, London, UK. ¹⁶Department of Psychiatry, University of Halle, Halle, Germany. ¹⁷Department of Biomedicine, Aarhus University, Denmark. ¹⁸Bioinformatics Research Centre (BiRC), Aarhus University, Aarhus, Denmark. ¹⁹Concentris Research Management GmbH, Fürstfeldbruck, Germany. ²⁰Leyden Delta B.V., Nijmegen, The Netherlands. ²¹Department of Psychiatry, Landspítali University Hospital, Reykjavik, Iceland. ²²Centre for Addiction and Mental Health, Toronto, Ontario, Canada. ²³Psychosis Research Unit, Aarhus University Hospital, Risskov, Denmark. ²⁴MRC Centre for Neuropsychiatric Genetics and Genomics, Institute of Psychological Medicine and Clinical Neurosciences, School of Medicine, Cardiff University, Cardiff, UK. ²⁵Division of Psychiatry Research, The Zucker Hillside Hospital, Northwell Health System, Glen Oaks, New York, USA. ²⁶Department of Genomics, Life & Brain Center, University of Bonn, Bonn, Germany. ²⁷Mental Health Services in the Capital Region of Denmark, Mental Health Center Copenhagen, University of Copenhagen, Denmark. ²⁸Department of Psychological Medicine, Institute of Psychiatry, Psychology and Neuroscience, King's College London, London, UK. ²⁹deCODE Genetics, Reykjavik, Iceland. ³⁰Center for Psychiatric Genomics, Department of Genetics, University of North Carolina, Chapel Hill, North Carolina, USA. ³¹Department of Medical Epidemiology and Biostatistics, Karolinska Institutet, Stockholm, Sweden.

GERADI Consortium Members

Denise Harold^{1,9}, Rebecca Sims¹, Amy Gerrish¹, Jade Chapman¹, Valentina Escott-Price¹, Richard Abraham¹, Paul Hollingworth¹, Jaspreet Pahwa¹, Nicola Denning¹, Charlene Thomas¹, Sarah Taylor¹, John Powell², Petroula Proitsi², Michelle Lupton², Simon Lovestone^{2,10}, Peter Passmore³, David Craig³, Bernadette McGuinness³, Janet Johnston³, Stephen Todd³, Wolfgang Maier⁴, Frank Jessen⁴, Reiner Heun⁴, Britta Schurmann^{4, 5}, Alfredo Ramirez⁴, Tim Becker⁶, Christine Herold⁶, André Lacour⁶, Dmitriy Drichel⁶, Markus Nothen⁷, Alison Goate⁸, Carlos Cruchaga⁸, Petra Nowotny⁸, John C. Morris⁸, Kevin Mayo⁸, Peter Holmans¹, Michael O'Donovan¹, Michael Owen¹ and Julie Williams¹.

¹Medical Research Council (MRC) Centre for Neuropsychiatric Genetics and Genomics, Neurosciences and Mental Health Research Institute, Department of Psychological Medicine and Neurology, School of Medicine, Cardiff University, Cardiff, UK. ²King's College London, Institute of Psychiatry, Department of Neuroscience, De Crespigny Park, Denmark Hill, London. ³Ageing Group, Centre for Public Health, School of Medicine, Dentistry and Biomedical Sciences, Queen's University Belfast, UK. ⁴Department of Psychiatry, University of Bonn, Sigmund-Freud-Straße 25, 53105 Bonn, Germany. ⁵Institute for Molecular Psychiatry, University of Bonn, Bonn, Germany. ⁶Deutsches Zentrum für Neurodegenerative Erkrankungen (DZNE), Bonn. ⁷Department of Genomics, Life & Brain Center, University of Bonn, Bonn, Germany. ⁸Departments of Psychiatry, Neurology and Genetics, Washington University School of Medicine, St Louis, MO 63110, US. ⁹Present address: Neuropsychiatric Genetics Group, Department of Psychiatry, Trinity Centre for Health Sciences, St James's Hospital, Dublin. ¹⁰Present address: Department of Psychiatry, University of Oxford.